



Politecnico di Torino

## Porto Institutional Repository

[Proceeding] Energy and water vapor transport in a turbulent stratified environment

*Original Citation:*

Gallana, L.; De Santi, F.; Iovieno, M.; Tordella, D. (2015). *Energy and water vapor transport in a turbulent stratified environment*. In: 68th American Physical Society - Division of Fluid Dynamics Annual Meeting 2015, Boston, 22 - 24 novembre 2015.

*Availability:*

This version is available at : <http://porto.polito.it/2630416/> since: February 2016

*Publisher:*

American Physical Society

*Terms of use:*

This article is made available under terms and conditions applicable to Open Access Policy Article ("Public - All rights reserved") , as described at [http://porto.polito.it/terms\\_and\\_conditions.html](http://porto.polito.it/terms_and_conditions.html)

Porto, the institutional repository of the Politecnico di Torino, is provided by the University Library and the IT-Services. The aim is to enable open access to all the world. Please [share with us](#) how this access benefits you. Your story matters.

(Article begins on next page)

Abstract Submitted  
for the DFD15 Meeting of  
The American Physical Society

**Energy and water vapor transport in a turbulent stratified environment** LUCA GALLANA, Politecnico di Torino, FRANCESCA DE SANTI, MICHELE IOVIENO, Politecnico di Torino, DIMEAS, RENZO RICHIARDONE, Universita' degli Studi di Torino, DANIELA TORDELLA, Politecnico di Torino, DIMEAS — We present direct numerical simulations about the transport of kinetic energy and unsaturated water vapor across a thin layer which separates two decaying turbulent flows with different energy. This interface lies in a shearless stratified environment modeled by means of Boussinesq's approximation. Water vapor is treated as a passive scalar (Kumar et al. 2014). Initial conditions have  $Fr^2$  between 0.64 and 64 (stable case) and between -3.2 and -19 (unstable case) and  $Re_\lambda = 250$ . Dry air is in the lower half of the domain and has a higher turbulent energy, seven times higher than the energy of moist air in the upper half. In the early stage of evolution, as long as  $|Fr^2| > 1$ , stratification plays a minor role and the flows follows closely neutral stratification mixing. As the buoyancy terms grows,  $Fr^2 \sim O(1)$ , the mixing process deeply changes. A stable stratification generates a separation layer which blocks the entrainment of dry air into the moist one, characterized by a relative increment of the turbulent dissipation rate compared to the local turbulent energy. On the contrary, an unstable stratification slightly enhances the entrainment. Growth-decay of energy and mixing layer thickness are discussed and compared with laboratory and numerical experiments.

Daniela Tordella  
Politecnico di Torino

Date submitted: 01 Aug 2015

Electronic form version 1.4